

PRODUCTION MODEL ANALYSIS ON ATLANTIC BIGEYE TUNA AS OF 1990

Pereira, J., Universidade dos Açores, 9900 Horta, Açores, Portugal

N. Miyabe

*National Research Institute of Far Seas Fisheries, Fishery Agency,
7-1, Orido 5 Chome, Shimizu-shi, 424 Japan*

SUMMARY

A production model analysis was conducted for the Atlantic stock of bigeye tuna, on the basis of CPUE from the Japanese longline fishery, for the years 1961-1990. The adjusted values of CPUE for 1974 have been used in the analysis. The fishing intensity estimated by the Honma method was adjusted for the deep longline operation, for 1980-1990. MSY estimated values are 67,500 MT for $m=2$, and 76,100 MT for $m=1$. Results show that catches in recent years are in the proximity of the estimated value of MSY. The analysis also indicates that current fishing effort is at a level below the F_{opt} estimated by the model.

RESUME

Cet article analyse l'état du stock de patudo atlantique par le modèle global, en utilisant les CPUE de la pêcherie palangrière japonaise pour les années 1961-90. Des valeurs révisées pour la CPUE de 1974 ont été utilisées. L'intensité de pêche a été estimée par la méthode de Honma en ajustant pour l'utilisation de la palangre profonde, pour la période 1980-90. Les valeurs estimées pour la PME sont de 67.500 TM pour $m = 2$, et de 76.100 TM pour $m = 1$. Les résultats montrent que les prises des années récentes sont à un niveau proche de ceux estimés pour la PME. L'analyse indique aussi que l'effort de pêche actuel est à niveau inférieur au F_{opt} estimé par le modèle.

RESUMEN

Este artículo analiza el estado del stock de patudo atlántico mediante el modelo global, utilizando las CPUE de la pesquería palangrera de Japón para los años 1961-1990. Se han utilizado los valores revisados para la CPUE de 1974. La intensidad de pesca ha sido estimada mediante el método de Honma ajustado a la utilización del palangre profundo, para el período 1980-90. Los valores estimados para el RMS son de 67.500 t para $m=2$ y de 76.100 t para $m=1$. Los resultados muestran que las capturas actuales se encuentran al nivel estimado para el RMS. El análisis indica también que el esfuerzo de pesca actual se encuentra a un nivel inferior al F_{opt} estimado por el modelo.

1. The bigeye tuna fishery in the Atlantic.

Bigeye tuna are widely distributed in the tropical and temperate waters of the Atlantic Ocean, between approximately 45° N and 45° S. Small bigeye have been observed in the only known nursery, which is located in the Gulf of Guinea.

The stock is exploited in the entire area of distribution by different fleets and gears, such as longline, purse seine and baitboats.

The main bigeye fishery is the longline, which operates throughout the year in the entire distribution area. The longline fishery directly targets large fish and, since 1980, Japanese and Korean longliners have targeted bigeye using deep longline and concentrating the effort in the time-area strata where the density of bigeye is higher.

Bigeye tuna is also targeted seasonally by some local baitboat fisheries in the areas of the Azores, Madeira and the Canary Islands. These fisheries of the Northeastern Atlantic islands exploit mainly medium and adult bigeye tuna. Dakar based baitboats, operating off Senegal and Mauritania, also target seasonally on medium size bigeye.

The eastern tropical surface fleets also catch an important amount of juvenile bigeye tuna, in mixed schools with young yellowfin and skipjack. Those fisheries do not target directly on bigeye tuna, but the catches in number of small bigeye are important.

Figure 1 shows the historical catch series for the Atlantic bigeye tuna stock, from 1961 up to 1990, by major fishing gears. In recent years the catch increased to a maximum of 74.400 MT in 1985, then decreased in the following years until 1987. In 1990 the catch level increased up to 69.300 MT.

2. Materials and methods

It is presently accepted the hypothesis that the bigeye tuna forms a unique stock in the Atlantic Ocean. The available information on tagging experiments, distribution by areas and sizes and spawning activities has shown evidence to support the hypothesis, which is adopted in the present study.

The Japanese longline fishery dominates in the Atlantic bigeye catch and targets this species over the entire area of distribution. The absence of an abundance index from the surface fisheries, qualified the use of Japanese statistics as a base of production model analysis to access the status of stock of this species.

The catch data are taken from the most recent ICCAT statistical publications (ICCAT, 1991) and those available at the ICCAT secretariat. The procedure used to estimate the total effective effort and catch of the Atlantic bigeye tuna is described by Miyabe (1988) and outlined as follows.

- Step 1. The Japanese longline Task II data were used to compute the effective effort of the species, converting the nominal fishing effort of deep longline operation to that of conventional longline operation on the basis of gear efficiency and deployment rate of deep longline, as described by Kume (1985). Tables 1 and 2 and Figures 2 and 3, give the relevant information mentioned in this step.
- Step 2. The Japanese nominal fishing effort thus calculated forms a base to compute effective effort by Honma method (Honma, 1973). Average year period to obtain average density index by month and 5° x 5° squares, covers the years from 1967 to 1972. The area south of 45° N and north of 40° S, except Mediterranean was chosen for the whole Atlantic for the computation of the effective effort.
- Step 3. The effective effort on bigeye by the Japanese longline fleet was raised to the total effective effort of all fishing gears multiplying the ratio of the total catch to the Japanese catch.

The method described in Step 3, was not used in the calculation of the total Atlantic effort for the year 1974. For this particular year, the cpue value used was an average of those observed in 1973 and 1975. In fact, the 1974 cpue value deduced from the Japanese statistics, appears unrealistic high which induces a bias in the estimation of bigeye abundance. A detailed analysis of bigeye

abundance in the period 1973-75 is presented by Pereira (1990).

A production model analysis was conducted using the program "PRODFIT" (Fox, 1975), for four cases of the shape parameter m , $m=0$, $m=1$, $m=2$ and m variable. The k parameter, corresponding to the number of year classes that mainly contributes to the fishery, was chosen as 4.

The input data used for Prodfit estimations are listed in table 3.

The present study updates the previous one by Pereira and Miyabe (1991), and covers the period from 1961 to 1990.

3. Results and discussions

3.1. CPUE trends

The annual trends of bigeye CPUE for the Japanese longline fishery, calculated based on standardised effort and catch in number of fish (catch in number per 100 hooks) is shown in figure 4. This longline CPUE is a good index of abundance of the adult stock, since the catch of the longline fishery is composed of medium and large size fish. This trend indicates that the CPUE has been relatively stable in the recent period, with a slightly increasing trend when compared to the period before the introduction of deep longline. Average CPUE for the last five years period (1986-90) is at 81.9% of the CPUE calculated for the beginning of the fishery, period 1961-1965.

3.2. Production model analysis

The results of the adjustment of the model to the input data, for each case of the m parameter are given in table 4 and in figure 5.

The MSY estimated values, depending on the value of the m parameter, are 67.500 MT for $m=2$, 76.100 MT for $m=1$ and 144.900 MT for $m=0$. In case of estimated m ($m=2.39$) the estimated MSY is 66.200 MT. The optimum effort levels estimated for each case range from 386 to 569 millions hooks, for $m=2$ and $m=1$ respectively. The estimated values for F_{opt} are higher than those observed in the fishery for recent years. This has been the case in previous analysis, where the

current fishing effort has been at a level lower than the optimal estimated by production model analysis.

The observed total catches of bigeye tuna in the Atlantic have decreased from 1986 to 1988, mainly due to a decline in the longline effort, but after 1989 the number of longliners in the Atlantic increased, and the MSY estimated value indicates that the present catches are at the level of the estimated MSY.

References

- FOX, W. 1975.- Fitting the generalised production model by least squares and equilibrium approximation. Fish. Bull. US. Vol. 73 (1): 23-36
- HONMA, M. 1973.- Overall fishing intensity and catch by length class of yellowfin tuna in Japanese Atlantic longline fishery, 1956-1971. ICCAT, Col. Vol. Sci. Papers, I: 59-77
- ICCAT, 1991.- Statistical bulletin, Vol. 21 - 1990.
- KUME, S. 1985.- An analysis on the stock abundance of Atlantic bigeye tuna caught by Japanese longline fishery. ICCAT, Col. Vol. Sci. Papers, XXIII (2): 248-253
- MIYABE, N. 1989.- An updated production model analysis on Atlantic bigeye tuna as of 1986. ICCAT, Col. Vol. Sci. Papers, XXX (1): 1-5
- PEREIRA, J. 1990.- Analyse de l'abondance de patudo Atlantique en 1973-75. ICCAT, Col. Vol. Sci. Papers, XXXII (1): 73-82
- PEREIRA, J. and MIYABE, N. 1991. Production model analysis on Atlantic bigeye tuna as of 1988. ICCAT, Col. Vol. Sci. Papers, XXXV (1): 22-27

Year		Area			
		1	2	3	4
1980	Rate (%)	5	41	75	29
	Efficiency	1.35	1.24	1.32	1.22
1981	Rate (%)	25	49	68	22
	Efficiency	1.78	1.51	1.11	1.21
1982	Rate (%)	29	67	85	38
	Efficiency	1.31	1.14	1.19	0.95
1983	Rate (%)	11	68	82	37
	Efficiency	1.75	1.25	1.16	1.07
1984	Rate (%)	10	63	90	50
	Efficiency	1.59	1.58	1.16	0.98
1985	Rate (%)	12	83	95	59
	Efficiency	1.82	1.62	1.17	0.86
1986	Rate (%)	8	91	96	35
	Efficiency	3.11	1.53	1.72	1.38
1987	Rate (%)	14	93	98	40
	Efficiency	1.59	1.27	1.48	0.89
1988	Rate (%)	8	96	96	64
	Efficiency	1.24	1.76	1.11	0.85
1989	Rate (%)	10	95	97	74
	Efficiency	2.39	0.98	1.02	1.40
1990	Rate (%)	6	90	95	60
	Efficiency	1.44	0.90	1.26	0.69

Table 1. Annual deployment rate of deep longline operation and gear efficiency factor over conventional longline for 1980 - 90.

Year	Catch in number (1,000)	Yield in Weight (1,000 MT)	Effective effort (million)	Hook rate
1961	243.7	11.0	27.3	0.892
1962	367.9	15.7	51.7	0.712
1963	285.3	14.5	45.8	0.623
1964	343.7	17.3	58.0	0.593
1965	648.3	28.5	113.8	0.570
1966	232.1	17.6	46.8	0.496
1967	180.9	8.5	30.5	0.593
1968	204.6	10.3	30.0	0.682
1969	263.3	10.3	36.8	0.717
1970	187.3	9.0	32.4	0.578
1971	394.9	20.3	83.0	0.476
1972	346.0	18.1	77.6	0.446
1973	391.3	20.0	74.7	0.524
1974	457.4	20.9	66.9	0.683
1975	449.1	17.4	110.4	0.407
1976	171.0	7.3	49.2	0.348
1977	190.0	9.2	35.2	0.540
1978	209.2	9.3	47.3	0.442
1979	270.4	12.0	62.2	0.435
1980	451.7	20.5	101.4	0.446
1981	469.2	21.0	130.8	0.359
1982	699.8	32.9	144.0	0.486
1983	351.6	15.1	77.7	0.452
1984	524.3	24.3	108.3	0.484
1985	674.8	31.6	119.4	0.565
1986	499.3	22.8	115.4	0.433
1987	404.5	18.6	73.9	0.547
1988	681.0	31.7	107.6	0.631
1989	860.0	39.4	145.2	0.592
1990	757.0	35.0	131.9	0.574

Table 2. Catch, fishing effort and CPUE on Atlantic bigeye tuna for the Japanese longline fishery, 1961-1990. For 1980-1990, deep longline efficiency was adjusted to the conventional gear.

Year	Yield in Weight (1,000 MT)	Effective effort (million)	CPUE
1961	17.0	42.2	0.403
1962	23.0	75.7	0.304
1963	25.9	81.8	0.317
1964	23.3	78.1	0.298
1965	39.2	156.5	0.250
1966	24.9	66.2	0.376
1967	24.7	88.6	0.279
1968	23.0	67.0	0.343
1969	35.8	127.9	0.282
1970	41.3	148.7	0.276
1971	54.9	224.5	0.245
1972	46.5	199.4	0.233
1973	56.5	211.0	0.268
1974	63.8	299.1	0.213
1975	60.6	384.5	0.158
1976	44.6	300.6	0.148
1977	54.2	207.4	0.261
1978	51.6	262.4	0.197
1979	45.1	233.8	0.193
1980	62.8	310.6	0.202
1981	67.2	417.9	0.161
1982	72.8	318.6	0.228
1983	58.5	301.0	0.194
1984	68.6	305.7	0.224
1985	74.4	281.1	0.265
1986	58.9	298.1	0.198
1987	48.7	193.5	0.252
1988	57.9	196.5	0.295
1989	69.0	254.3	0.271
1990	69.3	261.1	0.265

Table 3. Catch, fishing effort and CPUE on bigeye tuna for the total Atlantic, 1961-1990.

m (shape parameter)	Degree of fit index	Y - max (1,000 MT)	F - optimum (million hooks)	1985 - 90 catch (1,000 MT)
0	.5292	144.9	∞	48.7 - 74.4
1	.9290	76.1	569.1	
2	.9713	67.5	386.1	
variable (m=2.39)	.5388	66.2	359.2	

Table 4. Estimated population parameters obtained from production model analysis for Atlantic bigeye tuna, 1961 - 90.




Figure 1. Atlantic bigeye tuna catches (cumulative), by surface and longline gears, 1961 - 1990.




Figure 2. Area division used for the adjustment of deep longline effort..




Figure 4. Trend of annual CPUE of bigeye tuna caught by Japanese longline fishery in the Atlantic Ocean, 1961-1990.




Figure 3. Efficiency factor of deep longline operation over conventional longline, in the four areas used for the deep longline adjustment, years 1980-1990.




Figure 5. Yield curves of the production model analysis for bigeye tuna in the Atlantic, 1961 - 1990.