

SWORDFISH GROWTH PATTERN IN THE STRAIT OF GIBRALTAR; IMPLICATIONS FOR MIXING AMONG ATLANTIC AND MEDITERRANEAN STOCKS

Noureddine Abid¹, George Tserpes², M'Hamed Idrissi³ and Mohammed Bakkali⁴

SUMMARY

A growth study of the swordfish in the Strait of Gibraltar was carried out, based on monthly size frequencies data collected from the Moroccan driftnet fishery during the period 2006-2011. The growth parameters were estimated by the modal progression analysis (MPA), using both the Bhattacharya and NORMSEP methods. The standard von Bertalanffy growth function (VBGF) for length was found to be: $L_t = 253.6 [1 - \exp(-0.17(t + 1.30))]$. The growth pattern of swordfish in the Strait of Gibraltar was found to be very similar to that obtained from past studies in various Mediterranean areas. Given the existing growth differences among Atlantic and Mediterranean swordfish, this suggests that the majority of fish caught in this area are most likely belonging to the Mediterranean stock. However, further studies are needed to identify the degree of mixing among stocks.

RÉSUMÉ

Une étude de croissance de l'espadon dans le détroit de Gibraltar a été menée, sur la base des données mensuelles des fréquences de taille recueillies auprès de la pêcherie marocaine de filets dérivants pendant la période 2006-2011. Les paramètres de croissance ont été estimés par l'analyse de progression modale (MPA), en utilisant les méthodes Bhattacharya et NORMSEP. La fonction de croissance standard de von Bertalanffy (VBGF) pour la longueur s'est avérée être : $L_t = 253.6 [1 - \exp(-0.17(t + 1.30))]$. Le schéma de croissance de l'espadon dans le détroit de Gibraltar s'est avéré être très similaire à celui obtenu d'études antérieures dans diverses zones de la Méditerranée. Compte tenu des différences de croissance entre l'espadon de l'Atlantique et celui de la Méditerranée, ceci suggère que la majorité des poissons capturés dans cette zone appartiennent très vraisemblablement au stock méditerranéen. Il est toutefois nécessaire de procéder à de nouvelles études afin d'identifier le degré de mélange des stocks.

RESUMEN

Se llevó a cabo un estudio del crecimiento del pez espada en el estrecho de Gibraltar, basado en datos de frecuencias de talla mensuales recopilados en la pesquería de redes de enmalle marroquí durante 2006-2011. Los parámetros de crecimiento se estimaron por medio de un análisis de progresión modal (MPA), utilizando los métodos Bhattacharya y NORMSEP. Se halló que la función de crecimiento estándar de von Bertalanffy (VBGF) para la talla era: $L_t = 253,6 [1 - \exp(-0,17(t + 1,30))]$. Se descubrió que el patrón de crecimiento del pez espada en el estrecho de Gibraltar era muy similar al obtenido en pasados estudios en diversas zonas del Mediterráneo. Dadas las diferencias existentes de crecimiento entre el pez espada del Atlántico y del Mediterráneo, esto sugiere que la mayoría de los peces capturados en esta zona es más probable que pertenezcan al stock del Mediterráneo. Sin embargo, son necesarios más estudios para identificar el grado de mezcla entre los stocks.

KEYWORDS

Swordfish, Growth, Strait of Gibraltar, Stocks mixing

¹ National Institute for fisheries research (INRH), regional centre of Tangier. P.O. Box 5268, Tangier, Morocco

² Hellenic Centre for Marine Research, P.O. Box 2214, 71003, Heraklion, Greece

³ International Commission for the Conservation of Atlantic Tunas (ICCAT), Corazón de María

⁴ Faculté des Sciences et Techniques de Tanger. B.P. 416.Tanger, Maroc

1. Introduction

Swordfish is a cosmopolitan species found in the tropical and temperate waters of all the oceans, between 45°N and 45°S. In the Atlantic, the distribution includes the Mediterranean Sea, the Black Sea and the Marmara Sea (Palko *et al.*, 1981).

In the Mediterranean, swordfish spawn mainly around the Balearic Islands, in the center and south of the Tyrrhenian Sea, in the Ionian Sea, in the Strait of Messina and in the Levant basin (Beardsley, 1978; Rey, 1988; Tserpes *et al.*, 2001). In the northwest Atlantic, swordfish spawn all year, mainly in the Gulf of Mexico, south of the Sargasso Sea, east of the Antilles, in the Strait of Florida and along the southeast coast of the United States (Beardsley, 1978; Rey, 1988; Arocha and Lee, 1996).

Research results based on genetic studies have demonstrated that Mediterranean swordfish compose a unique stock separated from the north and south Atlantic stocks. Although, there is incomplete information on stock mixing and boundaries, it is generally believed that stock mixing is low and limited to the region around the Straits of Gibraltar (Anon., 2012).

The annual catches of the Mediterranean swordfish have fluctuated between 12,000-16,000 t over the last 15 years. With regard to the north Atlantic stock, the annual catches averaged about 11,551 t in the last decade. The Mediterranean swordfish is considered overfished (Anon., 2011), while the North Atlantic stock has been recently rebuilt to the B_{MSY} target, following a long-term management plan established by ICCAT since 1999 (Anon., 2010).

The growth parameters are essential elements in population dynamic models used by the International Commission for the conservation of Atlantic tunas (ICCAT) for the swordfish stocks assessment. Detailed information and update of these parameters is necessary to take into account growth variability among areas, as well as, changes due to biological and environmental factors.

The age and growth of swordfish have been mostly studied using anal fin spine sections (Berkeley & Houde, 1983; Tsimenides & Tserpes, 1989; Megalofonou *et al.*, 1990; Tserpes and Tsimenides, 1995; Ehrhardt, 1996; Aliçli and Oray., 2001). Few authors, however, have determined the age of the species either from otoliths (Radtke & Hurley, 1983; Wilson & Dean, 1983), or using length-frequency data (Ovchinnikov *et al.*, 1980; De Metrio & Megalofonou, 1987; El Hananch, 1987). Beckett (1974) studied the growth of the Atlantic swordfish from vertebrae.

In the Strait of Gibraltar, a mixing area between the Mediterranean and North Atlantic swordfish stocks, very little information is available on the biology and particularly on the growth of the species. In the present work we attempt to estimate the swordfish growth pattern in the area based on length frequency data, as the captured animals are directly exported to the European market making difficult to obtain hard parts for aging purposes.

Given that the aforementioned past studies have clearly demonstrated the existence of growth differences among Atlantic and Mediterranean stocks, our results would help to determine the stock mixing levels in the straits of Gibraltar. It should be noted that accomplishment of studies that can provide information on swordfish stock structure are highly recommended by ICCAT (Anon., 2012), which is the responsible body for the management of large pelagic stocks in the Atlantic and the Mediterranean.

2. Materials and Methods

During the period from 2006 to 2011, the size sampling of swordfish was conducted regularly at the Tangier port, located in the Strait of Gibraltar. All the sampled fish came from the Strait of Gibraltar and the Atlantic adjacent, the mixing area between the Mediterranean and the North Atlantic swordfish stocks (**Figure 1**).

A total of 23,979 swordfish (both sexes combined) were sampled for size and/or weight from Moroccan driftnet fishery operating in this area; from which 3,287 paired observations of length and weight were collected for estimating the monthly length-weight relationships.

The size of fish was measured to the nearest centimeter from the tip of the lower jaw to the fork of the tail (Lower Jaw Fork Length, LJFL), using a measuring tape. The size data were aggregated by 5cm intervals to estimate the monthly size frequencies.

The modal progression analysis (MPA) was applied to the monthly length frequency distributions by year. This analysis involves three stages:

- (i) decomposition of composite distributions into their components to identify means representing different age groups, using the Bhattacharya's Method (1967). The estimated mean lengths were then used as initial guesses to predict the mean length of the identified age groups, using the NORMSEP method based on the maximum likelihood concept to separation of the normally distributed components of size frequencies distributions (Hasselblad, 1966; Abrahamson, 1971; Pauly & Caddy, 1985).
- (ii) Subjective identification and linking of the means perceived to belong to the same cohorts and
- (iii) Using the growth increment data from the linking to estimate growth parameters.

The growth parameters were estimated by fitting the Standard VBGF (1) to the growth increment data using 2 different methods.

$$L_t = L_{00} (1 - e^{-k(t-t_0)}) \quad (1)$$

- *Fabens method:*

Fabens (1965) suggested a method for estimating L_{00} and K , by predicting length at recapture (L_r) based on the current parameter selection and the length at marking (L_m). The growth parameters are estimated by minimizing the sum of squares of errors (SSE), ie. The squared differences between the observed length at second reading (L_r) and the predicted length (L_{ri}):

$$SSE = \sum (L_{ri} - L_{ri}')^2$$

- *Appeldoorn's method:*

Appeldoorn (1987) and Soriano & Pauly (1989) suggested a method allowing the use of growth increment data to estimate the parameters of a seasonally oscillating version of the VBGF. They used Marquardt's algorithm for a nonlinear fit to minimize the following function:

$$SSE = \sum \{L_{i+\Delta t} - (L_{00} - L_i) e^{-(K\Delta t - S_t + S_{t+\Delta t})}\}^2$$

Where:

$$S_t = (CK/2\pi) \cdot \sin(2\pi(t - t_s))$$

$$S_{t+\Delta t} = (CK/2\pi) \cdot \sin(2\pi((t + \Delta t) - t_s))$$

And

$$t_s = 0.5 + WP$$

C: degree of oscillation

In this analysis, C and WP were set to 0 in order to estimate the parameters of the Standard VBGF that doesn't take into account the seasonal oscillation of growth.

The estimation of t_0 is based on the Von Bertalanffy plot, using the linear regression:

$$Y = a + bt$$

$$Y = -\ln(1 - (L_t/L_{00}))$$

t: arbitrary age of each cohort

L_t : mean length of each cohort at the time t

L_{00} : asymptotic length

$$t_0 = -a/b$$

The arbitrary age of each cohort was calculated after assigning an age group to the first mean length of each identified cohort, based on the length-age keys available from the literature (Berkely & Houde, 1983; Megalofonou *et al.*, 1990; Tserpes & Tsimenedes, 1995; Aliçli and Oray, 2001; Arocha *et al.*, 2003). The 1st July was assumed as the birth date of different cohorts (Megalofonou *et al.*, 1990; Tsimenedes & Tserpes 1989)

3. Results

Table 1 summarizes the number of fish sampled by month and year during the study period. Most of fish (34%) was sampled during the month of May corresponding to the peak of fishing activity of the Moroccan driftnet fleet in the Strait of Gibraltar. The size data are missing for certain months especially from September to December of the period 2006-2008. This could be explained by the fact that there was no size sampling conducted during that period because of the low swordfish landing at the Tangier port.

The monthly size distributions of swordfish during the whole period study were polymodal, which means that the catches are composed of many age groups. The size of fish ranges from 81 to 251 cm LJFL, but sizes between 110 and 190 cm represent roughly more than 85% of the catches (**Figure 2**).

The length-weight relationship for the whole fishing season is illustrated in **Figure 3**. The estimated length-weight relationship parameters (a, b), the number of fish sampled, the size range, the coefficient of determination (R^2) corresponding to the length weight-relationships, as well as the F test results, for the whole season and by month, are summarized in **Table 2**. All the F test results are statistically highly significant at 1% level, indicating that the slopes of the relationships are quite different from 0 ($b > 0$).

The Student test results are statistically highly significant at 1% level, which means that the parameter b is greater than 3, indicating that swordfish has a positive allometry. This means that its growth rate in weight is higher than that in length (**Table 3**).

The F test results of the analysis of variance from the General Linear Modeling Approach (GLM) and month as a factor showed that the length-weight relationships slopes differ statistically among months ($F = 11.51$, $p < 0.001$) (**Table 4**). It is noticeable that for a given size, the fish reached its highest weight in May. This indicates that the fish has a higher condition factor (fat fish) in that month which coincides with intense spawning activities (**Figure 4**).

Table 5 shows the mean size of different age groups identified from Bhattacharya and NORMSEP methods. Up to five (5) age groups were successfully decomposed from the monthly size distributions. **Table 6** displays the progression of the mean size of the cohorts: 2004, 2005, 2006, 2007 and 2008, from July 2006 to August 2011, based on the **Table 3**.

Given the observed growth rate of swordfish, the time interval chosen was large enough to easily detect the progression of the mean size of the different cohorts. The evolution of mean size of the cohorts from July 2006 to August 2011 is illustrated in **Figure 5**.

The current study suggests that swordfish caught in Strait of Gibraltar grows annually between 34cm for young fish (98 cm) and 9 cm for old fish (195 cm) (**Table 7**).

Figure 6 displays the mean size of the synthetic cohort aged from 14 to 85 months old (1 to 7 years old). It should be noted that for a given age between 21 and 52 months, the differences in the mean sizes between the different cohorts are less than 5%. This suggests that fish of the same age group have similar growth rates, and there are negligible cohort effects.

The growth parameters estimated from the current study are summarized in **Table 8**. Both the Appeldoorn and Fabens models fit well to the data ($R^2 = 0.63$) (**Figure 7**) and the estimated growth parameters are quite similar. The Von Bertalanffy plot for estimating the age at zero length (t_0) is displayed in **Figure 8**.

4. Discussions

The growth parameters estimated by the current study are similar to those found by other authors for the Mediterranean swordfish (**Table 9**) and consequently, the same is valid for the predicted mean sizes at age, which are similar to those obtained from previous studies in the Mediterranean (Megalofonou *et al.*, 1990; Tserpes & Tsimenides, 1995; Aliçli and Oray, 2001). Our size-at-age estimates are particularly close to those reported by Tserpes & Tsimenides (1995), except for the 0-age group (**Table 10 and Figure 9**). This difference is probably due to the fact that 0-age fish are not present in our samples. Consequently, the model didn't fit well the mean size for this age group. In general, the size at age differences among the current study and the above-mentioned ones are not greater than 6%.

In contrast, the predicted mean sizes at age from the current study are quite different from those estimated for the Atlantic swordfish (Arocha *et al.*, 2003), particularly for ages 1-5. In this case differences vary from 6 to 22%.

Our results are also in line with previous findings suggesting that the growth of the Mediterranean swordfish is rapid during the first 3 years of life, and then it slows down rapidly (Tserpes and Tsimenides, 1995; Aliçli and Oray, 2001). The fish grows about 32 cm during its first year of life, and then the annual growth increment decreases progressively to reach around to 10 cm at its 7th year of life.

The above findings indicate that the swordfish individuals caught in the Strait of Gibraltar belong more likely to the Mediterranean stock as it was already suggested by past genetic studies (Vinas *et al.*, 2007). The existence of mixing among Atlantic and Mediterranean swordfish in the area is well documented and it is believed that individuals migrate from the Atlantic to the Mediterranean for spawning (El Hannach, 1987; De la Serna and Alot, 1990; Abid, 1998). Although the higher condition factor in our data has been observed in May, which is a month of intense swordfish spawning activity in the Mediterranean, it seems that catches of Atlantic animals are not frequent enough to affect the growth parameter estimates at least for the young fish caught in this area. Given, however, that growth rate differences among stocks are relatively small at older ages, it is normal to expect that growth studies, such as the current one, will not be able to provide a complete picture of mixing in the Straits of Gibraltar.

As the results obtained from the statistical method are in agreement with those from the direct aging of the species using anal fine spine sections, the MPA used in this study can be considered an alternative approach to assess the growth of swordfish when hard parts are not available. However, this method presents some difficulties especially in decomposing size frequencies data into age groups for older fish (LJ-FL>200 cm). Further research considering sexually dimorphic growth may help to refine those questions.

References

- Abid, N. 1998. *Contribution à l'étude de la pêche marocaine de l'espadon dans le détroit de Gibraltar*. Mémoire de troisième cycle pour l'obtention du diplôme d'ingénieur d'État, spécialité : Halieutique. I.A.V, Hassan II. Rabat. 92 p
- Abrahamson, N.J. 1971. Computer programs for fish stock assessment. FAO. Fish. Tech. Pap. 101. pag.var.
- Anon., 2010. Report of the 2009 Atlantic swordfish Stock Assessment Session. Col. Vol. Sci. Pap. ICCAT, 65(1): 1-123.
- Anon., 2011. Report of the 2010 ICCAT Mediterranean Swordfish Assessment Meeting. Col. Vol. Sci. Pap. ICCAT, 66(4): 1405-1470.
- Anon., 2012. Report of the Standing Committee of Research and statistics (SCRS). ICCAT, report for biennial period, 2010-11, Part II (2011) - Vol. 2.
- Appeldoorn, R. 1987. Modification of a seasonally oscillating growth function for use with mark-recapture data. J.Cons. CIEM, 43: 194-198.
- Aliçli, T. Z., Oray, I. K., 2001. Age and growth (*Xiphias gladius* L., 1758) in the eastern Mediterranean Sea. Col. Vol. Sci. Pap. ICCAT, 52(2): 698-707.
- Arocha, F., Lee, D. W., 1996. Maturity at size, reproductive seasonality, spawning frequency, fecundity and sex ratio in swordfish from the Northwest Atlantic. Col. Vol. Sci. Pap. ICCAT, 45(2): 350-357.
- Arocha, F., Moreno, C., Beerkircher, L., Lee, D. W., Marciano, L., 2003. Update on the growth estimates for the swordfish, *Xiphias gladius*, in the Northwestern Atlantic. Col. Vol. Sci. Pap. ICCAT, 55(4): 1416-1429.
- Beardsley, J. L., 1978. Report of the swordfish workshop held at the Miami laboratory southeast fisheries center, National Marine Fisheries Service Miami Florida. June 7-9. Col. Vol. Sci. Pap. ICCAT, 7(1): 149-158.
- Beckett, J. S., 1974. Biology of swordfish, *Xiphias gladius* L., in the northwest Atlantic Ocean. Proceedings of the international Billfish symposium, Kailua-kona, Hawaii, 9-12 August 1972. Part 2. Review and contributed papers. pp: 103-106. July 1974.

- Berkeley, S. A., Houde, E.D., 1983. Age determination of broadbill swordfish, *Xiphias gladius*, from the Straits of Florida, using anal fin spine sections. U.S. Dep. Comm., NOAA, Tech. Rep. NMFS, 8: 137-143.
- Bhattacharya, C.G., 1967. A simple method of resolution of a distribution into Gaussian components. *Biometrics*, 23: 115-135.
- de la Serna, J. M., Alot, E., 1990. Consideraciones relativas a los desplazamientos efectuados por el pez espada (*Xiphias gladius*) en el área del estrecho de Gibraltar y otras observaciones relacionadas con la biología de la reproducción. *Col. Vol. Sci. Pap. ICCAT* 32(2): 353-359.
- Ehrhardt, N. M., 1996. On the age and growth of swordfish in the Northwest Atlantic Ocean. *Col. Vol. Sci. Pap. ICCAT*, 44 (2): 84-88.
- El Hannach, A., 1987. Données biologiques et écologiques sur l'espadaon (*Xiphias gladius*) L.1758 à partir de la pêche marocaine dans le détroit de Gibraltar. Thèse de Doctorat, spécialité halieutique. ENSA, Rennes, France. 162 p.
- Fabens, A.J., 1965. Properties and fitting of the von Bertalanffy growth curve. *Growth*, 29: 265-289.
- De Metrio, G., Megalofonou, P., 1987. Catch, size distribution, growth and sex ratio of swordfish (*Xiphias gladius* L) in the Gulf of Taranto. *FAO Fisheries report* 394: 91-102.
- Hasselblad, V., 1966. Estimation of parameters for a mixture of normal distributions. *Technometrics*, 8: 431-444
- ICCAT. 2003. Report for Biennial Period, 2002-2003, Part I. 207 pp.
- Megalofonou, P., Dean, J. M., De Metrio, G., 1990. First results on the aging of juvenile swordfish, *Xiphias gladius* L., from the Mediterranean Sea, using otoliths. *Col. Vol. Sci. Pap. ICCAT*, 33: 163-167.
- Ovchinnikov, V.V., Grudtsev, M. E., Kholodkova, S. V., 1980. Length-age composition of tropical Atlantic swordfishes (*Xiphias gladius*, L). *Col. Vol. Sci. Pap. ICCAT*, 9(3): 620- 623.
- Palko, B. J., Beardsley, G. L., Richards, W. J., 1981. Synopsis of the biology of the swordfish, *Xiphias gladius* Linnaeus. NOAA Technical Report NMFS Circular 441/FAO Fisheries Synopsis No. 127.
- Pauly, D., Caddy, J F., 1985. A modification of Bhattacharya's method for the analysis of mixtures of normal distributions. *FAO Fish. Circ.* (781): 16p.
- Radtke, R. L., Hurley, P. C. F., 1983. Age estimation and growth of broadbill swordfish, *Xiphias gladius*, from the NW Atlantic based on external features of otoliths. *US. Dep. Commer., NOAA tech. Rep. NMFS* 8: 145-150
- Rey, J. C., 1988. Comentarios sobre las areas de reproducción del pez espada, (*Xiphias gladius*) en el Atlántico y Mediterráneo. *ICCAT, Col. Vol. Sci. Pap.* 27(1): 180-193.
- Soriano, M.L., Pauly, D., 1989. A method for estimating the parameters of a seasonally oscillating growth curve from growth increment data. *ICLARM Fishbyte*, 7(1):18-21
- Tserpes, G., Tsimenides, N., 1995. Determination of age and growth of swordfish, *Xiphias gladius* L.1758, in the eastern Mediterranean using anal-end spines. *Fish. Bull.* 93: 594-602.
- Tserpes, G., Peristeraki, P., Somarakis, S., 2001. On the reproduction of swordfish (*Xiphias gladius* L) in the eastern Mediterranean. *Col. Vol. Sci. Pap. ICCAT*, 52(2): 740-744.
- Tsimenides, N., Tserpes, G., 1989. Age determination and growth of swordfish, *Xiphias gladius* L., 1785 in the Aegean Sea. *Fish. Res.*, 8(1989): 159-168.
- Vinas, J., Bremer, J. A., Mejuto, J., de la Serna, J. M., Garcias-Cortes, B., Pla, C., 2007. Swordfish genetic population structure in the north Atlantic and Mediterranean. *Col. Vol. Sci. Pap. ICCAT*, 61(1): 99-106.
- Wilson, C.A., Dean, J. M., 1983. The potential use of sagittae for estimating age of Atlantic swordfish, *Xiphias gladius*. *US. Dep. Commer. NOAA. Tech. Rep. NMFS* 8: 151-156

Table1. Number of fish sampled for size and or weight (LJFL) by year and by month during the period 2006-2011.

<i>Year/month</i>	<i>Apr.</i>	<i>May</i>	<i>Jun.</i>	<i>Jul.</i>	<i>Aug.</i>	<i>Sep.</i>	<i>Oct.</i>	<i>Nov.</i>	<i>Dec.</i>	<i>Total</i>
2006	406	438	165	283	273					1565
2007	92	527			278					897
2008	10	535	407		267					1219
2009		1442	691	261	259	203	287	760	449	4352
2010	190	161	158	307	162	211		905	315	2409
2011	2645	5173	1595	893	1955	633	168	69	406	13537
Total	3343	8276	3016	1744	3194	1047	455	1734	1170	23979

Table 2.Length weight relationship parameters (a and b), number, the size range, the coefficient of determination (R^2) as well as the F test results of AOV, by month and for the whole season for the period 2006-2010.

<i>Temporal strata</i>	<i>a</i>	<i>b</i>	<i>n</i>	<i>LJ-FL</i> <i>(cm)</i>	<i>R2</i>	<i>df</i>	<i>F</i>	<i>Pr (>F)</i>
Apr.	9×10^{-7}	3.541	692	90-214	0.95	1, 690	13856	< 2.2e-16
May	2×10^{-6}	3.411	1787	89-240	0.94	1,1785	26267	< 2.2e-16
Jun.	1×10^{-6}	3.529	320	95-234	0.95	1, 318	6721	< 2.2e-16
Jul.	2×10^{-6}	3.412	120	87-194	0.95	1,118	2227	< 2.2e-16
Aug.	6×10^{-6}	3.122	323	101-250	0.95	1,321	6343	< 2.2e-16
Oct.	7×10^{-7}	3.564	45	107-230	0.91	1,43	416	< 2.2e-16
Whole season	2×10^{-6}	3.373	3287	87-250	0.94	1,3285	52258	< 2.2e-16

Table 3. Student test results for the slope of the length-weight relationship (null hypothesis, $b=3$) of swordfish, by month and for the whole season, period 2006-2010.

<i>Temporal strata</i>	<i>n</i>	<i>Estimate</i>	<i>Standard error</i>	<i>t value</i>	<i>(Pr > t)</i>
Apr.	692	3.541	0.03008	17.985	<0.01
May	1787	3.411	0.02105	19.525	<0.01
Jun.	320	3.529	0.04305	12.288	<0.01
Jul.	120	3.412	0.07231	5.698	<0.01
Aug.	323	3.122	0.03918	3.114	<0.01
Oct.	45	3.564	0.1747	3.228	<0.01
Whole season	3287	3.373	0.01476	25.271	<0.01

Table 4. F test results of anova from GLM modeling

<i>Factor</i>	<i>df</i>	<i>Deviance</i>	<i>Resid. Df</i>	<i>Resid. Dev</i>	<i>F</i>	<i>Pr(> F)</i>
NULL			3286	1331.14		
Log.LJ.FL	1	1252.41	3285	78.73	57448.529	< 2.2e-16 ***
Month	5	6.08	3280	72.65	55.741	< 2.2e-16 ***
Log.LJ.FL:Month	5	1.25	3275	71.40	11.512	4.856e-11 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Table 5. Decomposition of monthly size distributions into age groups and their corresponding mean size using Bhattacharya and Normsep methods.

Date/age group	C1	C2	C3	C4	C5
Jul.06	110	131	172	208	
Aug.06	87	125	173	203	
Apr.07	110	132	162	202	
May.08	105	131	155	175	192
Jun.09	111	128	131	178	241
Oct.09	100	123	155	179	204
May.10	112	134	155	185	226
Nov.10	99	118	152	192	222
Aug.11	108	134	156	171	199

Table 6. Progression of the mean sizes and their standard deviation (Sd) of the 2004, 2005, 2006, 2007 and 2008 cohorts.

Date/cohort	2008		2007		2006		2005		2004	
	Mean size	Sd	Mean size	Sd	Mean size	Sd	Mean size	Sd	Mean size	Sd
Jul.06									110	9.4
Aug.06							87	4.6		
Apr.07							110	6.0	132	11.3
May.08					105	3.64	131	10.4	155	11.5
Jun.09			111	7.0	128	2.50			178	13.3
Oct.09	100	2.8	123	13.8			155	11.76		
May.10	112	6.4	134	6.2						
Nov.10					152	21.28			192	3.28
Aug.11	134	11.65	156	3.36					199	11.0

Table 7. Growth increment by size estimated for the 2004, 2005, 2006, 2007 and 2008 cohorts.

Date/cohort	2008		2007		2006		2005		2004	
	$\Delta L/\Delta T$	Mean size								
Apr.07							34	98	28	121
May.08							20	120	21	143
Jun.09					22	117			21	166
Oct.09			27	117			17	143		
May.10	20	106	20	128						
Nov.10					17	140			10	185
Aug.11	18	123	18	145					9	195

Table 8. Growth parameters estimated for swordfish from Gibraltar Strait based on growth increment data

Method	L_{00}	kK	t_0	R^2
Appeldoorn(1987)	253.6	0.170	-1.30	0.63
Fabens'(1965)	249.4	0.172	-1.23	0.62

Table 9. Growth parameters estimated for swordfish by several authors in different areas.

<i>Author</i>	<i>Method</i>	<i>Area</i>	<i>FL(cm)</i>	<i>L₀₀</i>	<i>K</i>	<i>T₀</i>
Berkeley & Houde (1983)	Spine	North Atlantic	??????	271cm	0.090	-3.79
Tserpes & Tsimenides (1995)	Spine	Eastern Mediterranean	62-210cm	238.6cm	0.185	-1.40
Aliçli (2000)	Spine	Eastern Mediterranean	52-219cm	252.2cm	0.133	-2.43
Megalofonou <i>et al.</i> (1990)	Spine	Eastern Mediterranean	71-207	243.8cm	0.140	-2.6
Arocha <i>et al.</i> (2003)	Spine	Northwest Atlantic	63-262cm	464.5 cm	0.002	-

Table 10. Predicted mean length by age of swordfish from different studies in the Atlantic (A), Mediterranean (M) and Gibraltar Strait (G)

<i>Age</i>	<i>Berkely&Houde(1982) (A)</i>	<i>Arocha et al (2003) (A)</i>	<i>Tserpes&Tsemindes (1995) (M)</i>	<i>Megalofonou et al. (1990) (M)</i>	<i>Alicli(2000) (M)</i>	<i>Currentstudy (G)</i>
0			55	74	70	50
1	99	105	86	97	92	82
2	119	129	111	116	112	109
3	137	146	133	132	130	132
4	153	160	151	147	145	151
5	167	171	166	160	158	167
6	180	180	178	171	170	180
7	192	189	188	180	180	192
8	202	196	197	189	189	201
9		203	204	196	197	210
10		210	210	202	204	216
11		216				
12		221				
13		227				
14		232				
15		236				
16		241				




Figure 1. Fishing grounds (shaded area) of the Moroccan driftnet fleet targeting swordfish in the Strait of Gibraltar.




Figure 2. Monthly size distribution of swordfish landed at the port of Tanger during the period 2006-2011.




Figure 3. Length weight relationship of swordfish caught in the Strait of Gibraltar (all data combined).




Figure 4. Comparison of the monthly length weight relationships of swordfish caught in the Strait of Gibraltar.




Figure 5. Evolution of the mean size of 2004, 2005, 2006, 2007 and 2008 cohorts from July 2006 to August 2011 (vertical bars represent standard deviation) LJFL: Lower jaw-Fork length. 24 month, 13 month, 22 month, 23 month and 15 month refer to the first age identified of the cohorts.





Figure 6. Evolution of the mean size of the synthetic cohort from 14 to 85 months, obtained from the juxtaposition of the plots of the Fig. 5. LJ-FL: Lower jaw-Fork length.

A



B




Figure 7. Growth curves of swordfish from Appeldoorn (A) and Faben's methods (B). Noise points represent observed data.




Figure 8. Von Bertalanffy plot for estimating the theoretical age at 0 length ($t_0 = -a/b$).




Figure 9. Comparison of swordfish growth curves between the current study and those used by ICCAT for the assessment of the Mediterranean (Tserpes & Tsimenides (1995) and Atlantic (Arocha *et al.*, 2003) stocks. LJ-FL: Lower jaw-Fork length.