

**Natural mortality rates adopted by the SCRS for major species**

Spp.	Stock	M (per year) <sup>1</sup> by age											Ref.
		0	1	2	3	4	5	6	7	8	9	10	
ALB	North, South		0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3			ANON. (1990)
BET	Atl.	0.8	0.8	0.4	0.4	0.4	0.4	0.4	0.4				ANON. (1984)
BFT	East		0.49	0.24	0.24	0.24	0.24	0.20	0.175	0.15	0.125	0.10	ANON. (1997);
BFT	West		0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14	?
SKJ	Atl.	0.8	0.8	0.8	0.8								ANON. (1999)
SWO	North		0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	ANON. (1987)
SWO	Med		0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	ANON. (1996)
YFT	Atl.	0.8	0.8	0.6	0.6	0.6	0.6						ANON. (1984)

<sup>1</sup> Last age with an M value is usually a "plus group"

**REFERENCES**

- ANON. (1984) Meeting of the working group of juvenile tropical tunas. Col. Vol. Sci. Pap. ICCAT, 21 (1): 1-289.
- ANON. (1987) Report of the Swordfish Assessment Workshop. Col. Vol. Sci. Pap. ICCAT, 26 (2): 339-395.
- ANON. (1990). Report of the 1989 ICCAT Albacore Workshop. Col. Vol. Sci. Pap. ICCAT, 31: 73-210
- ANON. (1996) Report of the First meeting of the Ad Hoc GFCM-ICCAT Working group on Stocks of Large Pelagic Fishes in the Mediterranean. Col. Vol. Sci. Pap. ICCAT, 45(1): 1-60.
- ANON. (1997) Report of the 1996 ICCAT SCRS Bluefin Tuna Stock Assessment Session. Col. Vol. Sci. Pap. ICCAT, 46 (1): 1-186
- ANON. (1999) 1998 Detailed Report on Skipjack Tuna. Col. Vol. Sci. Pap. ICCAT, 49 (3): 123-158

Last update: 4 August, 2005