

# Tagging of Atlantic bluefin tuna (*Thunnus thynnus*) with pop-up satellite archival tags (PSAT) in Norway during 2022

**Keno Ferter, Otte Bjelland, Jan Hinriksson, Martin Wiech, Leif Nøttestad**

Institute of Marine Research, Bergen, Norway

Corresponding author and project leader: Keno Ferter, Keno@hi.no, phone: +47 94103552



**Bergen, Norway, 3<sup>rd</sup> November 2022**

**Final project report prepared for the International Commission for the Conservation of Atlantic Tunas (ICCAT) / Grand Bluefin Year Programme (GBYP) 2022 – Phase 12. GBYP e-tagging program 2022.**



This project is co-funded by  
the European Union



## **Executive summary**

Atlantic bluefin tunas (BFT) have reoccurred in increasing numbers along the coast of Norway during the last years. To study the behavior, migration and general ecology of BFT returning to their historical feeding grounds in Norwegian waters, the Institute of Marine Research in Norway continued its tagging program of BFT along the coast of Norway in collaboration with the International Commission for the Conservation of Atlantic Tunas (ICCAT) between the 31<sup>st</sup> of August and 29<sup>th</sup> of September 2022. Like in previous years, the major aims were to collect genetic samples of BFT and tag these with both pop-up satellite archival tags (PSATs) and conventional tags as far north as possible. Tagging was performed on-board a specially designed tagging vessel with an aluminum ramp to pull the fish on board. In total, eleven BFT ranging from 237 cm to 283 cm (CFL) in length were tagged with PSATs and conventional tags, and genetic samples were collected. Eight BFT were caught from the tagging vessel, and three individuals were transferred from collaborating recreational fishing boats to the tagging vessel. All fish were caught using rod-and-line and spreader bars as lures. The results of this project contribute to the understanding of the behavior, migration and ecology of this highly migratory species at its historical feeding grounds far north in the northeast Atlantic Ocean.

## 1. Introduction

Atlantic bluefin tunas (BFT) have returned to Norwegian waters in large numbers during the last decade (Nøttestad et al., 2020). To study the behavior, migration and ecology of BFT in Norwegian waters, the Institute of Marine Research (IMR) in Norway conducted electronic tagging programs as part of the Grand Bluefin Year Programme (GBYP) in 2018, 2020 and 2021 (Ferter et al., 2019; Ferter et al., 2020; Ferter et al., 2021).

Like in previous years, the International Commission for the Conservation of Atlantic Tunas (ICCAT) provided pop-up satellite tags (PSATs) to be deployed in the Mediterranean and North Atlantic Ocean targeting eastern stock individuals as part of ongoing CPCs national electronic tagging programs (ICCAT GBYP CIRCULAR #0433/2022). Through a Memorandum of Understanding (MoU), IMR agreed to deploy five PSATs provided by ICCAT as part of the GBYP 2022 Phase 12 e-tagging activities. In addition, IMR had 6 self-financed PSATs available, and the results of these additional tags will be shared with ICCAT GBYP. Although both fishing and tagging were planned to be mainly performed from a research vessel owned by IMR, this study greatly benefited from collaboration with recreational BFT fishers in terms of knowledge sharing and helping to catch BFT for electronic tagging.


To study the behavior of individuals feeding at the northernmost distribution limit of the species, the aim of this project was to extend previous tagging campaigns by collecting genetic samples of BFT and tag these fish with PSATs and conventional tags during their feeding period as far north as possible in Norwegian waters.

## 2. Materials and Methods

All procedures followed the handling and tagging protocols in 2021 (Ferter et al., 2021).

### *Study area and period*

The field study was conducted north of 60°N between Bergen and Ålesund in western Norway between the 31<sup>st</sup> of August and 29<sup>th</sup> of September 2022 (Figure 1). This area was chosen because of many observations of feeding BFT during the study period, and because most of both recreational and commercial catches were taken within this area during this year's season. Moreover, like in previous years, the aim of this study was to deploy the PSATs as far north as possible in Norwegian waters. A total number of 16 effective fishing days were conducted from IMR's tagging vessel. All experimental procedures were approved by the Norwegian Food Safety Authority (FOTS ID 29603) and the Norwegian Directorate of Fisheries.



**Figure 1:** Map of the study area. The red dots indicate the tagging and release locations of the eleven tagged BFT in this study. Numbers next to the triangles indicate fish ID.

### *Fishing equipment and methods*

Eight of the fish were caught from the research vessel, and three from collaborating recreational fishing boats (table 1) using fishing rods in the 130 lbs range, and reels of at least size 80. All fish were fought stand-up using a fighting belt (Black Magic XL wide Equalizer gimbal and harness system). Spreader bars were used as fishing method on all fishing days. To keep fighting time to a minimum, the boat was used to overtake the fish as fast as possible after the first initial run. This made it possible to pull up the head of the fish from straight above, hampering the swimming of the fish and cutting angling duration to a minimum.

### *Tagging protocol*

Once the BFT were close to the boat, a large, barbless hook with a rope was placed between the tip of the tongue and the lower jaw of the BFT. The fish was then towed behind the boat (in 2 - 3 knots) for up to 10 minutes. Afterwards, the fish was pulled into the boat and placed on a 6 cm thick foam mattress which was covered with a smooth tarpaulin. The eyes of the fish were then covered with a towel to minimize stress, and a hose with high-volume but low-pressure continuous seawater supply was placed into the mouth of the fish to irrigate the gills. The fish were tagged with one PSAT (MiniPat-348, Wildlife Computers, 365 days deployment duration, constant pressure release after three days) fitted with two monofilament anchors (one of them fitted to the base of the PSAT and the other one as a loop) and titanium darts next to the second dorsal fin. In addition, a conventional spaghetti tag was placed close to the second dorsal fin following the instructions in the ICCAT-GBYP tagging manual (Cort et al., 2010). During tagging, the curved fork length (CFL) of the fish was measured and a fin clip was taken for genetic analysis. The fin clip was stored in > 99.0 % ethanol at 4°C. The genetic samples were sent to AZTI for further analysis, and to be included in the GBYP tissue bank. After tagging, the fish was released on a thin rope which was pulled through the same hole as the large, barbless hook, and towed behind the boat (in 2 - 3 knots) for a short time until the fish showed strong swimming activity.

## **3. Results**

A total number of eleven BFT were tagged with both PSATs and conventional spaghetti tags during the 16 fishing days (table 1). All fish were tagged north of 60°N (figure 1, table 1). The size of the fish ranged from 237 cm to 283 cm (CFL). Eight fish were caught from the research

vessel and three fish (IDs 1, 3 and 11) were transferred from a recreational fishing boat to the research vessel. Fighting time ranged between 18 to 40 minutes (mean 24 minutes).

**Table 1:** Overview of the eleven BFT tagged along the coast of Norway in 2022.

Date	ID	Boat	Release time	Release position	CFL [cm]	Angling duration [min]	PSAT ID	Conventional
01/09/2022	1	Recreational boat	16:00	62.381033, 5.206567	250	22	21P2007*	BYP031164
05/09/2022	2	Tagging boat	18:30	61.090283, 4.352467	280	30	21P2006*	BYP031163
11/09/2022	3	Recreational boat	11:40	62.295533, 5.009317	228	40	21P2005*	BYP031161
11/09/2022	4	Tagging boat	13:50	62.347350, 5.001050	240	20	21P2010*	BYP031162
23/09/2022	5	Tagging boat	14:07	61.489050, 4.344917	250	25	21P2016*	BYP031168/BYP031170
23/09/2022	6	Tagging boat	17:40	61.583300, 4.341217	283	30	21P0046	BYP031169
24/09/2022	7	Tagging boat	12:35	62.098633, 4.873667	248	20	21P0047	BYP031173
24/09/2022	8	Tagging boat	13:15	62.098633, 4.873667	237	20	21P0042	BYP031183
24/09/2022	9	Tagging boat	16:20	62.180783, 4.763067	251	18	21P0048	BYP031165
28/09/2022	10	Tagging boat	13:15	61.408917, 4.340600	254	20	21P0053	BYP031171
29/09/2022	11	Recreational boat	16:20	62.566067, 5.320133	277	20	21P0045	BYP031973

\* ICCAT GBYP PSAT

To date, none of the tags has reported, indicating that they are still attached to the fish, and that the fish survived.

#### 4. Discussion and concluding remarks

The 2022 tagging campaign was a continuation of previous electronic tagging of BFT in Norwegian waters (Ferter et al., 2020; Ferter et al., 2021). However, the number of tagged fish was higher than in previous years, and in total 25 fish have been tagged with PSATs in Norwegian waters since 2020. All 2022 tags are still attached to the BFT, indicating that the improved tagging protocol, i.e. on-board tagging with double anchoring, is an effective method. This is backed up by the fact that several of the tags from the 2020 and 2021 tagging projects stayed on for an entire year of deployment (unpublished data). As all fish in this year's study were tagged north of 60°N, the results of this study will further increase our knowledge on the behavior and migration of BFT at the northernmost border of its distribution range, and fill important knowledge gaps (Nøttestad et al., 2017; Horton et al., 2020; Nøttestad et al., 2020).

Pulling the fish on board for tagging makes it possible to place the tag accurately as desired. Results from both 2020, 2021 and 2022 demonstrate high retention rates of the tag and high survival rates (see also Stokesbury et al., 2011). Towing the fish on a thin rope after tagging until the fish showed strong swimming activity was a further improvement compared to previous tagging campaigns. Overall, the experiences and knowledge gained from this year's study have laid the foundation for future successful electronic tagging studies of BFT in Norwegian waters. Further tagging efforts are required to further improve our understanding of the migration pattern and behavior of BFT in its northernmost distribution area.

## 5. Acknowledgements

This work has been carried out under the ICCAT Atlantic-Wide Research Programme for Bluefin Tuna (GBYP), which is funded by the European Union, several ICCAT CPCs, the ICCAT Secretariat, and other entities (see <https://www.iccat.int/gbyp/en/overview.asp>). The content of this paper does not necessarily reflect ICCAT's point of view or that of any of the other sponsors, who carry no responsibility. In addition, it does not indicate the Commission's future policy in this area. Further funding was provided by the Institute of Marine Research in Norway through the Norwegian Sea program (Project 15698 – “Satellittmerking av makrellstørje”). Five of the PSATs and the conventional tags, in addition to other tagging equipment, were provided by ICCAT-GBYP. The tagging pole to set the conventional tags was provided by Anthony Mendillo Jr.. The authors are grateful to all the voluntary anglers and commercial fishers that supported the project by reporting sightings of BFT and developing fishing methods. A special thanks to “Team Big 1”, “Team Bluefin Raiders” and “Team Maguro Blue” for providing three of the fish. The authors are also thankful to the staff of the Institute of Marine Research, especially Gunnar Didriksen for his work efforts in the field.

## 6. References

- Cort, J. L., Abascal, F., Belda, E., Bello, G., Deflorio, M., de la Serna, J. M., Estruch, V., et al. 2010. Tagging Manual for the Atlantic-Wide Research Programme on Bluefin Tuna (ICCAT-GBYP). 45 pp.
- Ferter, K., Bjelland, O., Hinriksson, J., and Nøttestad, L. 2020. Tagging of Atlantic bluefin tuna (*Thunnus thynnus*) with pop-up satellite archival tags (PSAT) in Norway during 2020. p. 9. The International Commission for the Conservation of Atlantic Tunas.
- Ferter, K., Bjelland, O., Hinriksson, J., and Nøttestad, L. 2021. Tagging of Atlantic bluefin tuna (*Thunnus thynnus*) with pop-up satellite archival tags (PSAT) in Norway during 2021. p. 8. The International Commission for the Conservation of Atlantic Tunas.
- Ferter, K., Tracey, S., Hinriksson, J., Bjelland, O., Onandia, I., and Nøttestad, L. 2019. Tagging of Atlantic bluefin tuna (*Thunnus thynnus*) with pop-up satellite archival tags (PSAT) in western Norway during 2018. p. 12. The International Commission for the Conservation of Atlantic Tunas.
- Horton, T. W., Block, B. A., Drumm, A., Hawkes, L. A., O’Cuaig, M., Maoiléidigh, N. Ó., O’Neill, R., et al. 2020. Tracking Atlantic bluefin tuna from foraging grounds off the west coast of Ireland. *ICES Journal of Marine Science*, 77: 2066-2077.
- Nøttestad, L., Boge, E., and Ferter, K. 2020. The comeback of Atlantic bluefin tuna (*Thunnus thynnus*) to Norwegian waters. *Fisheries Research*, 231: 105689.
- Nøttestad, L., Tangen, Ø., Utne, K. R., and Hamre, J. 2017. Utbredelse, fangst og forskning av makrellstørje (*Thynnus thynnus*) i norsk økonomisk sone (NØS). 35 pp.
- Stokesbury, M. J., Neilson, J. D., Susko, E., and Cooke, S. J. 2011. Estimating mortality of Atlantic bluefin tuna (*Thunnus thynnus*) in an experimental recreational catch-and-release fishery. *Biological Conservation*, 144: 2684-2691.